

What is claimed is:

1.

A method of lowering egg cholesterol levels of eggs from laying fowl, comprising:
administering to an egg laying fowl an egg cholesterol lowering effective amount of a
5 polycosanol.

2.

The method of claim 1 wherein the polycosanol is at least C₂₀ in carbon chain
length.
10

3.

The method of claim 1 wherein the amount of polycosanol administered is from
0.01 mg/kg bodyweight/day to 1 mg/kg of bodyweight/day.

15

4.

The method of claim 3 wherein the amount of polycosanol administered is from .05
mg/kg of bodyweight to 0.5 mg/kg of bodyweight of the fowl.

5.

20

The method of claim 1 wherein administration is by mixing with normally fed
poultry feed.

6.

The method of claim 1 wherein the polycosanol is administered in conjunction with a poultry feed containing an edible oil.

5

7.

The method of claim 6 wherein the feed containing edible oil is selected from the group consisting of soybean, corn, cannula, sunflower and rapeseed oils.

8.

10

The method of claim 1 wherein the polycosanol is mixed with a micellular mix of a plant sterol.

9.

The method of claim 1 wherein the polycosanol is mixed with a micellular mix of a plant sterol and a phospholipid.

15

10.

The method of claim 9 wherein the phospholipid is lecithin.

11.

- A method of lowering egg cholesterol of eggs from egg laying fowl, comprising:
- mixing an egg cholesterol lowering effective amount of polycosanol with normal poultry feed to provide a feed mix; and
- 5 feeding said feed mix to egg laying fowl to lower egg cholesterol levels of eggs produced by said fowl.